

Bijections Between Multiparking Functions, Dirichlet Configurations, and Descending R -Traversals

Dimitrije N. Kostić*

Department of Mathematics
Texas A&M University, College Station, TX 77843
dkostic@math.tamu.edu

Key words and phrases. parking functions, critical configurations, spanning forests
Mathematics Subject Classification. 05C30, 05C05

Abstract

There are several combinatorial objects that are known to be in bijection to the spanning trees of a graph G . These objects include G -parking functions, critical configurations of G , and descending traversals of G . In this paper, we extend the bijections to generalizations of all three objects.

1 Introduction

A *parking function* (of length n) is a sequence $(a_1, a_2, \dots, a_n)$ of nonnegative integers for which there exists a permutation $\pi \in S_n$ so that $a_{\pi(i)} < i$ for every $1 \leq i \leq n$. This concept was introduced (by way of an analogy to parking on a one-way street) in a 1966 paper by Konheim and Weiss [KoWe] on linear probing. In combinatorics, parking functions are most famous for being in bijection to the set of labelled trees on n vertices and several combinatorial proofs of this fact are known; see [FoRi] for an example. More recently, renewed interest in parking functions has spurred various generalizations. In [Stan], Stanley showed that k -parking functions are in bijection to the chambers of the extended Shi arrangements and in [Yan] Yan showed that k -parking functions are in bijection to sequences of rooted b -forests. In [StPi], Stanley and Pittman showed that the number of $\vec{x}$ -parking functions is determined by the volume of a certain polytope.

Let G be a loopless graph with vertex set $V(G) = \{1, 2, \dots, n\}$. (Unless otherwise stated, we do not assume connectedness.) In 2004 Postnikov and Shapiro [PoSh] proposed a new generalization of the notion of a parking function, equivalent to the following. Let $\mathbb{N}$ be the set

*Partially supported by NSF VIGRE grant # 9977354.

of nonnegative integers and let G be connected. If $i \in U \subseteq V(G)$, the *out-degree* of i in U is $\mathbb{O}_U(i) := \#\{j \in V(G) - U \mid i \text{ is adjacent to } j\}$. The *in-degree* of i in U is $\mathbb{I}_U(i) := \#\{j \in U \mid i \text{ is adjacent to } j\}$. A *G -parking function* is a function $f : V(G) \rightarrow \mathbb{N} \cup \{\infty\}$ such that for any $U \subseteq V(G) - \{1\}$ there exists a vertex $i \in U$ such that $0 \leq f(i) < \mathbb{O}_U(i)$. In [ChPy], a family of bijections between the set of G -parking functions and spanning trees of G was constructed, each bijection being determined by a *proper set of tree orders*.

Kostić and Yan [KoYa] generalized this work further. A *G -multiparking function* is a function $f : V(G) \rightarrow \mathbb{N} \cup \{\infty\}$ such that for any $U \subseteq V(G)$ there exists $i \in U$ with either **(A)** $f(i) = \infty$, or **(B)** $0 \leq f(i) < \mathbb{O}_U(i)$. Throughout this paper, we will refer to those vertices i with $f(i) = \infty$ as *roots* and those with $0 \leq f(i) < \mathbb{O}_U(i)$ as being *well-behaved in U* . Let $\mathcal{MP} = \mathcal{MP}_{R,G}$ denote the set of G -multiparking functions with root set R .

There is a subtle but important difference between this definition of a G -multiparking function and the one that appears in [KoYa]. In that paper, the minimal vertex in each component of G is required to be a root; here there is no such restriction. Note, however, that R cannot be empty; $V(G)$ cannot have a well-behaved vertex, so it must have a root.

This paper will construct bijections between G -multiparking functions and two other objects, Dirichlet configurations and descending R -traversals. In section 2, we provide an algorithm to verify whether a function is a G -multiparking function. In section 3 we introduce Dirichlet configurations and some basic facts about them. In section 4 we establish a bijection between G -multiparking functions and Dirichlet configurations on G . In section 5 we introduce descending R -traversals, define a certain partition of the set of these objects, and prove that they are in bijection to G -multiparking functions.

2 A Burning Algorithm for Multiparking Functions

We begin with a simple result on G -multiparking functions. It is proven in [KoYa], but we present the proof here for completeness.


Lemma 2.1. *Let $f : V(G) \rightarrow \mathbb{N} \cup \{\infty\}$ be a vertex function. Then, f is a G -multiparking function if and only if there exists a permutation $\pi \in S_n$ such that the vertex $\pi(i)$ is either a root of f or well-behaved in the set $U_i := V(G) - \{\pi(1), \pi(2), \dots, \pi(i-1)\}$.*

Proof. Suppose $\pi \in S_n$ satisfies the condition and let $U \subseteq V(G)$. Let l be the maximum index such that $U \subseteq U_l$. By maximality, $\pi(l) \in U$. Thus for $i = \pi(l)$, either i is a root or $0 \leq f(i) < \mathbb{O}_{U_l}(i) \leq \mathbb{O}_U(i)$. The other implication follows from the definition of a G -multiparking function. $\square$


This lemma suggests a simple algorithm for determining whether a function is a G -multiparking function. Let $U_0 = V(G)$. If there is a vertex $i_1 \in U_0$ that is a root (clearly, nothing in U_0 can be well-behaved), then let $U_1 := U_0 - \{i_1\}$. If there is a vertex $i_2 \in U_1$ which is a root or well-behaved, then let $U_2 := U_1 - \{i_2\}$. Lemma 2.1 implies that, when this process is continued, $U_n = \emptyset$ if and only if f is a G -multiparking function. This algorithm is

a generalization of the *burning algorithm*, which was originally developed in [Dhar] to study critical configurations.

Example 1. In several examples in this paper, the following graph Γ will be considered.



The following example illustrates the burning algorithm described above, proving that the illustrated function is a Γ -(multi)parking function. In the leftmost picture the circled vertices are in U_1 . In the next leftmost picture the circled vertices are in U_2 , and so forth.



3 Dirichlet Configurations

Now we introduce a generalization of a structure that appears in the literature in a variety of contexts and, with minor variations, is known as a critical configuration, a sandpile model, and a chip-firing game. Let R be a set of vertices containing at least one vertex from each component of G . A *configuration* μ on G (with *root set* R) is an integer-valued function on the vertex set for which $\mu(i) = -\infty$ if $i \in R$ and $0 \leq \mu(i) < \infty$ otherwise. A vertex i is said to be *ready* (in μ) if $\mu(i) \geq \deg(i)$. μ is *stable* if $0 \leq \mu(i) < \deg(i)$ for every $i \notin R$. An *avalanche* is a finite sequence $\alpha = (\mu_1, \mu_2, \dots, \mu_t)$ of configurations on G , where for each $1 \leq s < t$ there exists a vertex $i_s \in V(G)$ which is ready in μ_s and


$$\mu_{s+1}(i) = \begin{cases} \mu_s(i) - \deg(i) & \text{if } i = i_s \\ \mu_s(i) + e(i, i_s) & \text{if } i \neq i_s \end{cases}$$

where $e(i, i_s)$ is the number of edges between i and i_s . If we think of μ_s as keeping track of how many “chips” are stored at each vertex on the graph, then we transform μ_s into μ_{s+1} by sending a chip down each edge adjacent to i_s . This process is often called *firing* a vertex (hence the “chip-firing” terminology), and so one usually thinks of an avalanche as a sequence of vertex firings. Note that only vertices that are ready can be fired, and that the same vertex may be fired several times in succession if it has a large enough number of chips. We say that α *begins* at μ_1 , *ends* at μ_t , and *connects* these two configurations. We use the convention that if, in any avalanche, μ_1 is stable then every vertex in R is fired in some

arbitrary but fixed order and that this is the only situation in which roots are fired. Note that if a chip is sent to a root it disappears from the system; it follows from the connectedness of each component of G that, given any configuration, there is an avalanche leading to a stable configuration. μ is *recurrent* if there is an avalanche that begins and ends at μ . μ is *Dirichlet* if it is both stable and recurrent. Let $\mathcal{DC} = \mathcal{DC}_{R,G}$ denote the set of Dirichlet configurations on G with root set R .

Dirichlet configurations are usually called *critical configurations* when G is connected, and this case has been studied extensively (see, for example, [Bigg1]). Aspects of Dirichlet configurations were first examined in [ChEl], such as bounds on the number of vertex firings necessary to reach a stable configuration.

Example 2. The following example illustrates a critical configuration for Γ . Every vertex is labelled “ v_i/n ”, where v_i is the vertex label and n is the number of chips at that vertex at that configuration. The vertex about to be fired in each configuration is circled.



Note that in the second configuration in the above avalanche, we could have fired either v_2 or v_3 . If we had fired v_2 instead of v_3 , we would still have ended the avalanche on the configuration we started with.

It is not difficult to see that Dirichlet configurations exist on every graph (for instance, the configuration with $R = V(G)$) and that for every configuration there is an avalanche ending on a stable configuration. This is essentially because every component contains a root and therefore the total number of chips on the graph is nonincreasing after the root firings (if the first configuration is stable) in an avalanche. See Lemma 1 of [ChEl] for a detailed proof.

The following is a characterization of recurrent configurations. Let χ be the configuration

$$\chi(i) = \begin{cases} 0 & \text{if } i \in R \\ \sum_{r \in R} e(i, r) & \text{if } i \notin R \end{cases}$$

Proposition 3.1. *The configuration μ is Dirichlet $\iff$ it is stable and there is an avalanche connecting $(\mu + \chi)$ to μ , where $(\mu + \chi)(v) = \mu(v) + \chi(v)$.*

Proof. ($\Leftarrow$) The trivial avalanche (consisting of firing all the roots only) connects μ to $(\mu + \chi)$, and thus concatenating this avalanche with the avalanche connecting $(\mu + \chi)$ to μ shows that μ is recurrent. Since it is stable, μ is Dirichlet.

($\Rightarrow$) Given a Dirichlet μ , it is stable and recurrent. Thus there is an avalanche $(\mu, \omega_1, \omega_2, \dots, \omega_l, \mu)$. But since μ is stable, the roots are the only vertices that can be fired first. Thus, $\omega_k = (\mu + \chi)$ where k is the number of roots. Thus, $(\omega_k, \omega_{k+1}, \dots, \omega_l, \mu)$ is the necessary avalanche. $\square$

Cori and Rossin [CoRo] have a similar proof for the case when the graph is connected. The set of critical configurations of G is closely related to the set of G -parking functions; the most famous connection is that both sets are in bijection to the spanning trees of G . Here, however, we provide a bijection between G -multiparking functions and Dirichlet configurations on G that does not go through the set of spanning trees. To simplify the presentation, we will assume G has no multiple edges.

4 A Bijection Between Dirichlet Configurations and G -Multiparking Functions

Theorem 4.1. *Fix a root set R and let $\mathcal{MP} = \mathcal{MP}_{\mathcal{R},G}$ and $\mathcal{DC} = \mathcal{DC}_{\mathcal{R},G}$. Define $\Omega : \mathcal{MP} \rightarrow \mathcal{DC}$ by $\Omega(f) = \Omega_f$ where*

$$\Omega_f(i) = \begin{cases} -\infty & \text{if } i \in R. \\ \deg(i) - 1 - f(i) & \text{if } i \notin R. \end{cases}$$

Then Ω is a bijection, whose inverse $\Omega^{-1} : \mathcal{DC} \rightarrow \mathcal{MP}$, is given by $\Omega^{-1}(\mu) = \Omega_\mu^{-1}$ where

$$\Omega_\mu^{-1}(i) = \begin{cases} \infty & \text{if } i \in R. \\ \deg(i) - 1 - \mu(i) & \text{if } i \notin R. \end{cases}$$

Proof. Let f be any G -multiparking function. First we show that Ω_f is a Dirichlet configuration. This is trivial if $R = V(G)$, so assume $R \subset V(G)$. As $\Omega_f(i) < \deg(i)$ for every vertex i , Ω_f is stable. By proposition 3.1, finding an avalanche connecting $(\Omega_f + \chi)$ to Ω_f is enough to show that Ω_f is recurrent.

Note that $(\Omega_f + \chi)(i) = \deg(i) - 1 - f(i) + \chi(i)$ for every nonroot vertex i . Therefore, a vertex i in the configuration $(\Omega_f + \chi)$ is ready if and only if $\chi(i) > f(i)$. Since f is a G -multiparking function, the set of all non-root vertices must have a well-behaved vertex, say j , and this implies $\chi(j) > f(j)$. Hence, $(\Omega_f + \chi)$ is not stable.

Let $i_1, i_2, \dots, i_n$ be a burning sequence for f (in the sense of Lemma 2.1), with $i_1, i_2, \dots, i_k$ as the roots of f . We have just shown that there is a vertex that can be labelled i_{k+1} . It is enough to show that if the vertices $i_{k+1}, \dots, i_{l-1}$ can be fired, then i_l can be fired. Notice that for any $U \subseteq V(G)$, $\deg(i_l) = \mathbb{O}_U(i_l) + \mathbb{I}_U(i_l)$. So if $U = \{i_l, i_{l+1}, \dots, i_n\}$, then firing i_{k+1} through i_{l-1} sends exactly $\mathbb{O}_U(i_l) - \chi(i_l)$ chips to i_l . So, i_l will have at least $\deg(i_l) - 1 - f(i_l) + \mathbb{O}_U(i_l)$ chips. Since f is a G -multiparking function, $f(i_l) < \mathbb{O}_U(i_l)$, so $\deg(i_l) - 1 - f(i_l) + \mathbb{O}_U(i_l) = \deg(i_l) - 1 - (f(i_l) - \mathbb{O}_U(i_l)) \geq \deg(i_l)$, and thus i_l will be ready. Hence, every non-root vertex in an avalanche beginning with $(\Omega_f + \chi)$ must be fired, and the throwing-out sequence specified is also a sequence in which the vertices can be fired. (Note that although there may be several throwing-out sequences for f , they all yield the same final configuration.) Note that μ is a Dirichlet configuration if and only if a firing sequence exists, and this argument can be reversed to obtain a burning sequence, proving that this correspondence is surjective.

Finally, we must show that this sequence of firings beginning at $(\Omega_f + \chi)$ ends at Ω_f . If i is any vertex, it loses $\deg(i)$ chips when fired. As its neighbors are fired, i recovers exactly $\deg(i) - \chi(i)$ chips, since the roots are not fired. Thus, at the end of this avalanche, i has exactly $\deg(i) - 1 - f(i) + \chi(i) - \deg(i) + (\deg(i) - \chi(i)) = \deg(i) - 1 - f(i)$ chips, meaning that we end on the configuration Ω_f .

Finally, it is obvious that Ω^{-1} is the inverse of Ω . $\square$

This result strengthens earlier work by Biggs (see Lemma 3(ii) in [Bigg2]). This simple bijection also provides information on the natural poset orders on the sets of G -multiparking functions and Dirichlet configurations with a given root set. If f is a G -multiparking function, it is immediate from the definition that any vertex function which is less than or equal to f on each vertex is also a G -multiparking function. This determines a simple poset order on the G -multiparking functions. Analogously, if μ is a Dirichlet configuration then any other configuration which is stable and greater than or equal to μ on every vertex is also Dirichlet. Hence there is also a simple poset order on the Dirichlet configurations and the Hasse diagrams of these two posets are identical, except that one is upside-down.

Corollary 4.2. *If f and g are G -multiparking functions, then $f \leq g$ (in the G -multiparking function poset order described above) if and only if $\Omega_f \geq \Omega_g$ (in the Dirichlet configuration poset order described above).*

Theorem 4.1 also suggests a burning-type algorithm for verifying that a configuration is Dirichlet for a given graph.

Corollary 4.3. *A configuration μ on G is Dirichlet $\iff$ there exists a permutation $\pi \in S_n$ such that for every vertex i , either $\pi(i)$ is a root or $\deg(\pi(i)) > \mu(\pi(i)) \geq \mathbb{I}_{U_i}(\pi(i))$, where $U_i := V(G) - \{\pi(1), \dots, \pi(i-1)\}$.*

Proof. By theorem 4.1, μ is critical if and only if $f := \Omega^{-1}(\mu)$ is a G -multiparking function, and this is true if and only if there is a permutation $\pi \in S_n$ such that $0 \leq f(\pi(i)) < \mathbb{O}_{U_i}(\pi(i))$ for every nonroot vertex i . But this is true if and only if

$$\begin{aligned} & 0 \leq \deg(\pi(i)) - 1 - \mu(\pi(i)) < \mathbb{O}_{U_i}(\pi(i)) \\ \iff & \deg(\pi(i)) - 1 \geq \mu(\pi(i)) > \deg(\pi(i)) - 1 - \mathbb{O}_{U_i}(\pi(i)) \\ \iff & \deg(\pi(i)) > \mu(\pi(i)) \geq \mathbb{I}_{U_i}(\pi(i)) \end{aligned}$$

$\square$

We will hereafter refer to the permutations in proposition 4.3 as *Dirichlet certificates* for μ . This proposition also helps us identify the avalanches connecting a Dirichlet configuration to itself.

Proposition 4.4. *Let μ be a Dirichlet configuration and let $\pi \in S_n$. π is a Dirichlet certificate for $\mu \iff$ the avalanche determined by the firing sequence $\pi(1), \pi(2), \dots, \pi(n)$ connects μ to itself.*

Proof. ($\Leftarrow$) Let $\alpha = (\mu = \mu_1, \mu_2, \dots, \mu_n, \mu_1)$ be the avalanche determined by π and suppose $\pi(i)$ is a nonroot. We must show that $\deg(\pi(i)) > \mu_1(\pi(i)) \geq \mathbb{I}_{U_i}(\pi(i))$ for every such $i \leq n$. By assumption, μ is a Dirichlet configuration, so μ is stable, and thus $\deg(\pi(i)) > \mu_i(\pi(i))$ for every i .

The structure of α is that the vertices $\pi(1), \dots, \pi(i-1)$ are fired, and after these firings we arrive at μ_i . Then $\mu_i(\pi(i)) - \mu_1(\pi(i)) = \mathbb{O}_{U_i}(\pi(i))$, since $U_i = V(G) - \{\pi(1), \dots, \pi(i-1)\}$. Also, $\pi(i)$ is ready in μ_i and therefore $\mu_i(\pi(i)) \geq \deg(\pi(i))$. Thus

$$\begin{aligned} \mathbb{O}_{U_i}(\pi(i)) &= \mu_i(\pi(i)) - \mu_1(\pi(i)) \\ &\geq \deg(\pi(i)) - \mu_1(\pi(i)) \\ &= \mathbb{O}_{U_i}(\pi(i)) + \mathbb{I}_{U_i}(\pi(i)) - \mu_1(\pi(i)) \end{aligned}$$

Thus we have $\mu_1(\pi(i)) \geq \mathbb{I}_{U_i}(\pi(i))$, proving that π is a Dirichlet certificate for μ_1 .

($\Rightarrow$) If π is a Dirichlet certificate, then $\deg(\pi(i)) > \mu_1(\pi(i)) \geq \mathbb{I}_{U_i}(\pi(i))$ for every nonroot $\pi(i)$. Since $\mu = \mu_1$ is Dirichlet, it is stable, and thus only the roots can be fired. Suppose $\pi(1), \dots, \pi(i-1)$ have been fired in that order. Assuming $\pi(i)$ is not a root, $\mu_i(\pi(i)) = \mu_1(\pi(i)) + \mathbb{O}_{U_i}(\pi(i)) \geq \mathbb{I}_{U_i}(\pi(i)) + \mathbb{O}_{U_i}(\pi(i)) = \deg(\pi(i))$, and so $\pi(i)$ is ready in μ_i . Thus, $\pi(1), \dots, \pi(n)$ defines an avalanche.

It is clear that this avalanche connects μ_1 to itself, since we begin at that configuration and every vertex is fired exactly once, meaning that the net change in chips at each vertex i is $\sum_{j \neq i} e(i, j) - \deg(i) = 0$. $\square$

5 Descending Traversals

Let G be as above, but connected and with a total ordering $<_E$ on the edge set $E(G)$ and $V(G) = [n]$. Let $m = n + \#E(G)$. Let $\Sigma = \Sigma(G) = (\sigma_1, \sigma_2, \dots, \sigma_m)$ be a sequence of the edges and vertices of G in which each edge and vertex appears exactly once. Let $\Sigma^{\leq i} := (\sigma_1, \sigma_2, \dots, \sigma_i)$ and $\Sigma^{\geq i} := (\sigma_i, \sigma_{i+1}, \dots, \sigma_m)$. (Similarly, $\Sigma^{< i} := (\sigma_1, \sigma_2, \dots, \sigma_{i-1})$ and $\Sigma^{> i} := (\sigma_{i+1}, \sigma_{i+2}, \dots, \sigma_m)$.) We define Σ to be a *descending traversal on G* if it satisfies three conditions:

1. σ_1 is a vertex,
2. σ_i ($i \neq 1$) a vertex $\Rightarrow \sigma_{i-1}$ is an edge adjacent to σ_i ,
3. σ_i an edge $\Rightarrow$ it is adjacent to a vertex σ_k with $k < i$ and σ_i is maximal with respect to $<_E$ among all edges in $\Sigma^{\geq i}$ that are adjacent to some vertex in $\Sigma^{< i}$.

This definition is due to Cori and LeBorgne [CoLe]. They provided explicit bijections from the descending traversals to the spanning trees and from the descending traversals to the critical configurations, and hence a bijection between these other two objects.

Now assume G is the same as above, except not necessarily connected, and $R \subseteq V(G)$. Let Σ^* be a list of some vertices and edges of G (Σ^* contains no repetitions). Let $\mathbb{E}(\Sigma^*)$

be the set of edges not in Σ^* which are adjacent to a vertex in Σ^* . We let Π_i be the set of ordered pairs (Σ^*, W) where $W \subseteq \mathbb{E}(\Sigma^*)$ and where not both of $W = \emptyset$ and $R \subseteq \Sigma^*$ is true. A *choice function* on G is any function ζ from Π_i to $E(G) \cup R$ such that

$$\zeta(\Sigma^*, W) \begin{cases} \in W & \text{if } W \neq \emptyset \\ \in R - \Sigma^* & \text{if } W = \emptyset \text{ and } R \not\subseteq \Sigma^* \end{cases}$$


Fix a choice function ζ and let $\Sigma = (\sigma_i)_{i=1}^m$ be a sequence containing each edge and vertex of G exactly once. We call Σ a *descending R -traversal on G* , where $R = \{\sigma_{s_1}, \sigma_{s_2}, \dots, \sigma_{s_k}\}$, such that each subsequence $S_i = (\sigma_{s_i}, \sigma_{s_i+1}, \dots, \sigma_{s_{i+1}-1})$ of Σ satisfies:

1. $\sigma_{s_i} = \zeta(\Sigma^{<s_i}, \emptyset)$, where σ_{s_i} is a root,
2. $\sigma_j \in S_i, j > s_i$, a vertex $\Rightarrow \sigma_{j-1}$ is an edge adjacent to σ_j ,
3. $\sigma_j \in S_i$ an edge $\Rightarrow \sigma_j$ is adjacent to a vertex σ_k with $k < j$ and $\sigma_j = \zeta(\Sigma^{\leq j-1}, \mathbb{E}(\Sigma^{\leq j-1}))$.

Let $\mathcal{DT} = \mathcal{DT}_{R,G,\zeta}$ denote the set of descending R -traversals on G . Note that the first condition and the requirement that Σ can be partitioned into subsequences S_i is not very restrictive. To check that a subsequence is a descending R -traversal it is generally only necessary to confirm that the last two conditions hold.

If one defines $R := \{v_1\}$ and ζ to be the function that picks the largest-index edge available, then the descending R -traversals of G are, in fact, just the descending traversals of G .

Example 3. We illustrate some descending R -traversals of Γ , for different R . In all these examples, let $\zeta(\Sigma^*, W)$ be the largest-index edge in W if $W \neq \emptyset$ and the smallest vertex in R otherwise.



1. Let $R = \{v_1\}$. Then, $(v_1, e_4, v_2, e_3, e_2, v_4, e_5, v_3, e_1)$ and $(v_1, e_4, e_1, v_3, e_5, e_3, v_2, e_2, v_4)$ are descending R -traversals of Γ .
2. Let $R = \{v_2, v_3\}$. Then, $(v_2, e_4, e_3, e_2, v_3, e_5, e_1, v_1, v_4)$ and $(v_2, e_4, e_3, e_2, v_4, e_5, v_3, e_1, v_1)$ are descending R -traversals of Γ .
3. Let $R = \{v_1, v_2, v_4\}$. Then, $(v_1, e_4, e_1, v_2, e_3, v_3, e_5, e_2, v_4)$ is a descending R -traversal of Γ .

Now suppose Σ is a descending R -traversal, $R = \{\sigma_{s_1}, \sigma_{s_2}, \dots, \sigma_{s_k}\}$, and ζ is the choice function. With this as input, we define a function f_Σ on $V(G)$ in the following way:

Algorithm A

1. If $v = \sigma_{s_i}$ for some i , then set $f_\Sigma(v) = \infty$.
2. Otherwise, set $f_\Sigma(v) = j - 1$, where j is the number of edges adjacent to v that precede v in Σ .

Note that if $v \notin R$, then by condition (2) of the definition of a descending R -traversal, it is preceded by an edge adjacent to it. Thus, $f_\Sigma(v) = j - 1 \geq 0$ and so $f_\Sigma : V(G) \rightarrow \mathbb{N} \cup \{\infty\}$.

Proposition 5.1. $f_\Sigma \in \mathcal{MP}$ for any $\Sigma \in \mathcal{DT}$.

Proof. Let $f = f_\Sigma$ and let $\sigma_{v_1}, \sigma_{v_2}, \dots, \sigma_{v_n}$ be the vertex subsequence of Σ . We will show that this is a burning sequence for f , proving by Lemma 2.1 that f is a G -multiparking function. (It is clear that f has k roots.)

First, note that $f(\sigma_{v_1}) = \infty$. Let U_i be the set of vertices in $\Sigma^{\leq i}$. Now suppose $\sigma_{v_1}, \sigma_{v_2}, \dots, \sigma_{v_{i-1}}$ are all either roots or well-behaved in $U_1, U_2, \dots, U_{i-1}$, respectively. Suppose σ_{v_i} is not a root. If $f(\sigma_{v_i}) = j - 1$, then there are exactly j edges adjacent to σ_{v_i} and preceding it in Σ . Each of these edges is preceded in Σ by a vertex adjacent to it (note part 3 of the definition of a descending R -traversal). These vertices are among $\{\sigma_{v_1}, \sigma_{v_2}, \dots, \sigma_{v_{i-1}}\} = U_i$, and thus $0 \leq f(\sigma_{v_i}) = j - 1 < j = \mathbb{O}_{U_i}(\sigma_{v_i})$. Lemma 2.1 implies that $f \in \mathcal{MP}$. $\square$

Example 4. Let $\zeta(\Sigma^*, W)$ be the largest-index edge if $W \neq \emptyset$ and the lowest-index vertex in $R - \Sigma^*$ otherwise. Let $R = \{v_1, v_4\}$. Below is a table listing some descending R -traversals of Γ on the left-hand side and the corresponding (under algorithm A) Γ -multiparking functions on the right-hand side. (The list of descending R -traversals is not exhaustive.)

$(v_1, e_4, e_1, v_4, e_5, v_3, e_3, v_2, e_2)$	$\rightarrow$	$(\infty, 1, 1, \infty)$
$(v_1, e_4, e_1, v_4, e_5, v_3, e_3, e_2, v_2)$	$\rightarrow$	$(\infty, 2, 1, \infty)$
$(v_1, e_4, e_1, v_4, e_5, e_2, v_2, e_3, v_3)$	$\rightarrow$	$(\infty, 1, 2, \infty)$
$(v_1, e_4, v_2, e_3, e_2, v_4, e_5, e_1, v_3)$	$\}$	$(\infty, 0, 2, \infty)$
$(v_1, e_4, v_2, e_3, e_2, e_1, v_4, e_5, v_3)$		
$(v_1, e_4, e_1, v_3, e_5, v_4, e_3, v_2, e_2)$	$\}$	$(\infty, 1, 0, \infty)$
$(v_1, e_4, e_1, v_3, e_5, e_3, v_2, e_2, v_4)$		
$(v_1, e_4, e_1, v_3, e_5, v_4, e_3, e_2, v_2)$	$\}$	$(\infty, 2, 0, \infty)$
$(v_1, e_4, e_1, v_3, e_5, e_3, v_4, e_2, v_2)$		
$(v_1, e_4, v_2, e_3, e_2, v_4, e_5, v_3, e_1)$	$\}$	$(\infty, 0, 1, \infty)$
$(v_1, e_4, v_2, e_3, e_2, e_1, v_3, e_5, v_4)$		
$(v_1, e_4, v_2, e_3, v_3, e_5, e_2, e_1, v_4)$	$\}$	$(\infty, 0, 0, \infty)$
$(v_1, e_4, v_2, e_3, v_3, e_5, e_2, v_4, e_1)$		

Lemma 1 of [CoLe] states that if $(\sigma_i)_{i=1}^m$ and $(\tau_i)_{i=1}^m$ are descending traversals and k is the minimal index at which they differ, then one of σ_k and τ_k is an edge and the other is a vertex. The example above shows that this is not necessarily true for descending R -traversals; $(v_1, e_4, e_1, v_4, e_5, e_2, v_2, e_3, v_3)$ and $(v_1, e_4, e_1, v_3, e_5, v_4, e_3, e_2, v_2)$ do not observe this property.

Let $\Psi = \Psi_{R,G,\zeta} : \mathcal{DT} \rightarrow \mathcal{MP}$ be defined by $\Psi(\Sigma) = f_\Sigma$. The above example also illustrates that Ψ , as defined, is not generally injective. We will now define, for each graph G , root set R , and choice function ζ , a partition of $\mathcal{DT}$ over which Ψ will turn out to be injective.

Let f be any function from $V(G)$ to $\mathbb{N} \cup \{\infty\}$ such that $f(i) = \infty$ if and only if $i \in R$. We can consider $\Psi^{-1}(f)$, the (possibly empty) set of all descending R -traversals that are mapped to f . It is then clear that $\mathcal{R} = \mathcal{R}_{R,G,\zeta} := \{\Psi^{-1}(f) \mid \Psi^{-1}(f) \neq \emptyset\}$ is a partition of the set of descending R -traversals, where $\Psi^{-1}(f) = \{\Sigma \in \mathcal{DT} \mid \Psi(\Sigma) = f\}$. It is also clear that Ψ is constant over each $\Psi^{-1}(f)$ in $\mathcal{R}$, and that Ψ is injective when viewed as a function with $\mathcal{R}$ as its domain. Throughout the rest of this paper, we will view Ψ as a function from $\mathcal{R}$ to $\mathcal{MP}$.

Now we define an algorithm that will convert a a multiparking function to a descending R -traversal.

Algorithm B

- **Step 1: initial condition.** If $i = 1$ then $\Sigma^{\leq i} := (\zeta(\emptyset, \emptyset))$.
- **Step 2: insert the next entry.** Suppose $i > 1$. If there exists a vertex $v \notin \Sigma^{\leq i-1}$ such that $\Sigma^{\leq i-1}$ contains exactly $f(v) + 1$ edges adjacent to v , then $\Sigma^{\leq i} := \langle \Sigma^{\leq i-1}, v \rangle$. If no such vertex exists, then $\Sigma^{\leq i} := \langle \Sigma^{\leq i-1}, \zeta(\Sigma^{\leq i-1}, \mathbb{E}(\Sigma^{\leq i-1})) \rangle$. Repeat this step until $i = m$.

Example 5. Recall the conditions in Example 4. Below is a table listing all the Γ -multiparking functions on the right-hand side and the corresponding (under algorithm B) descending R -traversals of Γ on the right-hand side.

$(\infty, 1, 1, \infty)$	$\rightarrow$	$(v_1, e_4, e_1, v_4, e_5, v_3, e_3, v_2, e_2)$
$(\infty, 2, 1, \infty)$	$\rightarrow$	$(v_1, e_4, e_1, v_4, e_5, v_3, e_3, e_2, v_2)$
$(\infty, 1, 2, \infty)$	$\rightarrow$	$(v_1, e_4, e_1, v_4, e_5, e_2, v_2, e_3, v_3)$
$(\infty, 0, 2, \infty)$	$\rightarrow$	$(v_1, e_4, v_2, e_3, e_2, e_1, v_4, e_5, v_3)$
$(\infty, 1, 0, \infty)$	$\rightarrow$	$(v_1, e_4, e_1, v_3, e_5, e_3, v_2, e_2, v_4)$
$(\infty, 2, 0, \infty)$	$\rightarrow$	$(v_1, e_4, e_1, v_3, e_5, e_3, v_4, e_2, v_2)$
$(\infty, 0, 1, \infty)$	$\rightarrow$	$(v_1, e_4, v_2, e_3, e_2, e_1, v_3, e_5, v_4)$
$(\infty, 0, 0, \infty)$	$\rightarrow$	$(v_1, e_4, v_2, e_3, v_3, e_5, e_2, e_1, v_4)$

Proposition 5.2. $\Sigma^{\leq m} \in \mathcal{DT}$ for any $f \in \mathcal{MP}$.

Proof. We must first show that algorithm B can, in fact, always reach $\Sigma^{\leq m}$ if it acts on some $f \in \mathcal{MP}$. Clearly $\Sigma^{\leq 1}$ can be reached, so suppose $\Sigma^{\leq i} = (\sigma_j)_{j=1}^i$ can be reached for some $1 \leq i < m$. There are two cases in which Algorithm B might fail to reach $\Sigma^{\leq m}$.

First, suppose that there are two vertices v and w , neither in $\Sigma^{\leq i}$, such that there are exactly $f(v) + 1$ and $f(w) + 1$ edges in $\Sigma^{\leq i}$ that are adjacent to them respectively. (We may also assume, without loss of generality, that i is the minimum index at which there is more than one vertex ready to be appended to $\Sigma^{\leq i}$.) Note that the edge $\{v, w\}$, if it exists, is not in $\Sigma^{\leq i}$; no such edge could be in $\mathbb{E}(\Sigma^{\leq j})$ for any $j \leq i$ since neither v nor w is in $\Sigma^{\leq j}$. Therefore, in the sequence $\Sigma^{\leq i-1}, \Sigma^{\leq i-2}, \dots, \Sigma^{\leq 1}$ there must be a $\Sigma^{\leq j}$ which contains exactly $f(v) + 1$ edges adjacent to v but fewer than $f(w) + 1$ edges adjacent to w . Thus, v should have been added earlier and i does not exist.

We must also show that there is no index i for which $R \subseteq \Sigma^{\leq i}$ and $\mathbb{E}(\Sigma^{\leq i}) = \emptyset$. Let i be an index for which there is no vertex $v \notin \Sigma^{\leq i}$ adjacent to exactly $f(v) + 1$ edges in $\Sigma^{\leq i}$. Assume $R \subseteq \Sigma^{\leq i}$. Clearly, if $\Sigma^{\leq i}$ contains $V(G)$, then $\mathbb{E}(\Sigma^{\leq i})$ cannot be empty unless $i = m$. So let U be the set of vertices not in $\Sigma^{\leq i}$. Since $f \in \mathcal{MP}$ and there is no root in U , this set must have a well-behaved vertex. That is, there is a vertex $v \in U$ such that $0 \leq f(v) < \mathbb{O}_U(v)$. In particular, $0 < \mathbb{O}_U(v) \leq \#\mathbb{E}(\Sigma^{\leq i})$.

It is clear, from the construction of Algorithm B, that $\Sigma^{\leq m}$ satisfies the last two conditions in the definition of a descending R -traversal. $\square$

Let $\Phi = \Phi_{R,G,\zeta} : \mathcal{MP} \rightarrow \mathcal{DT}$ be defined by $\Phi(f) = \Sigma^{\leq m}$.

Proposition 5.3. Φ is injective.

Proof. Let f and g be different functions in $\mathcal{MP}$, and $\Phi(f) = \Sigma_f$ and $\Phi(g) = \Sigma_g$. Since f and g are different, there is a vertex v at which $f(v) < g(v)$ (v is not a root, since f and g have the same root set). There is an index i at which v appears in Σ_f . This means $\Sigma_f^{\leq i} = \langle \Sigma_f^{\leq i-1}, v \rangle$, but then either $\Sigma_f^{\leq i-1} \neq \Sigma_g^{\leq i-1}$ or $\Sigma_g^{\leq i} \neq \langle \Sigma_g^{\leq i-1}, v \rangle$ and so $\Phi(f) \neq \Phi(g)$. $\square$

Proposition 5.4. $\Psi(\Phi(f)) = f$ for any $f \in \mathcal{MP}$.

Proof. It is enough to show that $\Phi(f) \in \Psi^{-1}(f)$ for any $f \in \mathcal{MP}$. Suppose $\Phi(f) = (\sigma_i)_{i=1}^m$. Note that $\sigma_i \in R$ if and only if $\sigma_i = \zeta(\Sigma^{\leq i-1}, \emptyset)$ (where $\Sigma^{\leq i-1} = \emptyset$ if $i = 1$) and this is true if and only if $f(\sigma_i) = \infty$. Therefore $\Psi(\Phi(f))|_{\sigma_i} = \infty = f(\sigma_i)$. If $f(\sigma_i) = a$ for some vertex σ_i then $\Sigma^{\leq i-1}$ contains exactly $a + 1$ edges adjacent to σ_i . Therefore, $\Psi(\Phi(f))|_{\sigma_i} = a = f(\sigma_i)$. So, Ψ maps $\Phi(f)$ to f and thus $\Phi(f) \in \Psi^{-1}(f)$. $\square$

Acknowledgements

I extend my sincere gratitude to Professors Rob Ellis and Catherine Yan for their many helpful comments.

References

- [Bigg1] Biggs, Norman. *Chip Firing and the Critical Group of a Graph*. CDAM Research Report Series, 1995.
- [Bigg2] Biggs, Norman. *The Tutte Polynomial as a Growth Function*. Journal of Algebraic Combinatorics **10** (1999), 115-133.
- [ChPy] Chebikin, Denis and Pylyavskyy, Pavlo. *A Family of Bijections Between G -Parking Functions and Spanning Trees*. Journal of Combinatorial Theory A **110** no. 1 (2005), 31-41.
- [ChEl] Chung, Fan and Ellis, Rob. *A Chip-Firing Game and Dirichlet Eigenvalues*. Discrete Mathematics **257** (2002), 341-355.
- [CoLe] Cori, Robert and Le Borgne, Yvan. *The Sandpile Model and Tutte Polynomials*. Advances in Applied Mathematics **30** (2003), 44-52.
- [CoRo] Cori, Robert and Rossin, Dominique. *On the Sandpile Group of Dual Graphs*. European Journal of Combinatorics **21** (2000), 447-459.
- [Dhar] Dhar, D. *Self-Organized Critical State of the Sandpile Automaton Models*. Physical Review Letters **64** no. 14 (1990), 1613-1616.
- [DRSV] Dhar, D; Ruelle, P; Sen, S; and Verma, D-N. *Algebraic Aspects of Abelian Sandpile Models*. Journal of Physics A: Mathematical and General **28** (1995), 805-831.
- [FoRi] Foata, Dominique and Riordan, John. *Mappings of Acyclic and Parking Functions*. Aequationes Mathematicae **10** (1974), 10-22.
- [KoWe] Konheim, A.G. and Weiss, B. *An Occupancy Discipline and Applications*. SIAM Journal of Applied Mathematics **14** (1966), 1266-1274.
- [KoYa] Kostić, Dimitrije and Yan, Catherine. *Multiparking Functions, Graph Searches, and the Tutte Polynomial*. Preprint. See <http://arxiv.org/abs/math.CO/0607602>.
- [PoSh] Postnikov, Alexander and Shapiro, Bruce. *Trees, Parking Functions, Syzygies, and Deformations of Monomial Ideals*. Transactions of the American Mathematical Society **356** (2004), 3109-3142.
- [Stan] Stanley, Richard P. *Hyperplane Arrangements, Parking Functions, and Tree Inversions*. Appears in *Mathematical Essays in Honor of Gian-Carlo Rota*, B. Sagan and R. Stanley, eds. Birkhäuser, Boston/Basel/Berlin, 1988, 359-375.
- [StPi] Stanley, Richard P. and Pittman, Jim. *A Polytope Related to Empirical Distributions, Plane Trees, Parking Functions, and the Associahedron*. Discrete and Computational Geometry, **27** (2002), 603-634.

- [Yan] Yan, Catherine H. *Generalized Parking Functions, Tree Inversions, and Multicolored Graphs*. Advances in Applied Mathematics **27** (2001), 641-670.